

# Adora2a-KO

**Nomenclature** C57BL/6Smoc-*Adora2a*<sup>em1Smoc</sup>

**Cat. NO.** NM-KO-200018

**Strain State** Repository Live

## Gene Summary

|                               |                       |                                    |
|-------------------------------|-----------------------|------------------------------------|
| <b>Gene Symbol</b><br>Adora2a | <b>Synonyms</b>       | A2aR; A2AAR; AA2AR                 |
|                               | <b>NCBI ID</b>        | <a href="#">11540</a>              |
|                               | <b>MGI ID</b>         | <a href="#">99402</a>              |
|                               | <b>Ensembl ID</b>     | <a href="#">ENSMUSG00000020178</a> |
|                               | <b>Human Ortholog</b> | ADORA2A                            |

## Model Description

Exon 3 of Adora2a gene was deleted to generate Adora2a knockout mice.

**Research Application:** Therapeutic antibody target


\*Literature published using this strain should indicate: Adora2a-KO mice (Cat. NO. NM-KO-200018) were purchased from Shanghai Model Organisms Center, Inc..

## Disease Connection

|               |                     |   |
|---------------|---------------------|---|
| <b>Myopia</b> | <b>Phenotype(s)</b> | <a href="#">MGI:3622794</a>   |
|               | <b>Reference(s)</b> | Zhou X, Huang Q, An J, Lu R, Qin X, Jiang L, Li Y, Wang J, Chen J, Qu J, Genetic deletion of the adenosine A2A receptor confers postnatal development of relative myopia in mice. Invest Ophthalmol Vis Sci. 2010 Sep;51(9):4362-70 |

|                                       |                     |   |
|---------------------------------------|---------------------|---|
| <b>Primary Pulmonary Hypertension</b> | <b>Phenotype(s)</b> | <a href="#">MGI:5473699</a>   |
|                                       | <b>Reference(s)</b> | Xu MH, Gong YS, Su MS, Dai ZY, Dai SS, Bao SZ, Li N, Zheng RY, He JC, Chen JF, Wang XT, Absence of the adenosine A2A receptor confers pulmonary arterial hypertension and increased pulmonary vascular remodeling in mice. <i>J Vasc Res.</i> 2011;48(2):171-83 |

## Validation Data



**Fig1. Detection of Adora2a-ko expression in thymus and spleen by RT-PCR.** Abbr. M, DNA marker; HO, homozygous; HE, heterozygous; WT, wild type.